



**WOODROW FIRST SCHOOL & NURSERY**

LONGDON CLOSE  
WOODROW SOUTH  
REDDITCH  
WORCS  
B98 7UZ

[www.woodrowfirstschool.co.uk](http://www.woodrowfirstschool.co.uk)

**HEADTEACHER – MR R KIERAN  
HINTON**

**DEPUTY HEAD – MISS L**

TEL NO: 01527 527619

7 December 2022

Dear Parents & Carers

We have been advised of one confirmed case of Strep A and other children are being tested. We have been in contact with the Health Protection Agency for further guidance.

**School remains open.**

Regrettably, it **may** mean we will have to cancel some Christmas festivities:

- The children's Christmas singing and the Christmas Fayre etc

We will let you know as soon as we know more.

As ever, our aim is always to keep our community and children as safe as possible.

If you have any concerns about your children please see attached information or contact 111 for further guidance.

Yours sincerely

Mr R Kieran  
Headteacher



Dear Parent(s) / Carer(s),

**Re: Increase in scarlet fever**

We are writing to inform you of a recent [national/local] increase in notifications of scarlet fever to the UK Health Security Agency (UKHSA), above seasonal expected levels.

We would like to take this opportunity to remind you of the signs, symptoms and the actions to be taken if you think that you or your child might have scarlet fever.

**Signs and symptoms of scarlet fever**

Scarlet fever is a common childhood infection caused by *Streptococcus pyogenes*, or group A Streptococcus (GAS). It is not usually serious, but should be treated with antibiotics to reduce the risk of complications (such as pneumonia) and spread to others. The early symptoms of scarlet fever include sore throat, headache, fever, nausea and vomiting. After 12 to 48 hours, the characteristic red, pinhead rash develops, typically first appearing on the chest and stomach, then rapidly spreading to other parts of the body, and giving the skin a sandpaper-like texture. The scarlet rash may be harder to spot on darker skin, although the 'sandpaper' feel should be present. Patients typically have flushed cheeks and be pale around the mouth. This may be accompanied by a bright red red 'strawberry' tongue.

If you think you, or your child, might have scarlet fever:

- contact your GP or NHS 111 as soon as possible
- make sure that you or your child take(s) the full course of any antibiotics prescribed. Although you or your child will feel better soon after starting the course of antibiotics, you must complete the course to ensure that you do not carry the bacteria in your throat after you have recovered
- stay at home, away from nursery, school or work for at least 24 hours after starting the antibiotic treatment, to avoid spreading the infection

You can help stop the spread of infection through frequent hand washing and by not sharing eating utensils, clothes, bedding and towels. All contaminated tissues should be disposed of immediately.

**Invasive Group A Strep (iGAS)**

The same bacteria which cause scarlet fever can also cause a range of other types of infection such as skin infections (impetigo) and sore throat. In very rare cases, the bacteria can get into the bloodstream and cause an illness called invasive group A strep (iGAS). Whilst still very uncommon, there has been an increase in iGAS cases

this year, particularly in children under 10 years old. It is very rare for children with scarlet fever to develop iGAS infection.

As a parent, you should trust your own judgement.

Contact NHS 111 or your GP if:

- your child is getting worse
- your child is feeding or eating much less than normal
- your child has had a dry nappy for 12 hours or more or shows other signs of dehydration
- your baby is under 3 months and has a temperature of 38C, or is older than 3 months and has a temperature of 39C or higher
- your baby feels hotter than usual when you touch their back or chest, or feels sweaty
- your child is very tired or irritable

Call 999 or go to A&E if:

- your child is having difficulty breathing – you may notice grunting noises or their tummy sucking under their ribs
- there are pauses when your child breathes
- your child's skin, tongue or lips are blue
- your child is floppy and will not wake up or stay awake

### **Stop the spread**

During periods of high incidence of scarlet fever, there may also be an increase in outbreaks in schools, nurseries and other childcare settings. Children and adults with suspected scarlet fever should stay off nursery / school / work until **24 hours** after the start of appropriate antibiotic treatment. Good hygiene practice such as hand washing remains the most important step in preventing and controlling spread of infection.

### **Resources**

[NHS – Scarlet Fever](#)

[Scarlet fever: symptoms, diagnosis and treatment](#)

[Health protection in education and childcare settings](#)

[Hand hygiene resources for schools](#)